

Wastewater usage in urban and peri-urban agricultural production systems: scenarios from India

PREM JOSE VAZHACHARICKAL ^{*a} and SUMITA GUPTA GANGOPADHYAY ^{bc}

** Corresponding Author, Email: -premjosev@gmail.com*

a. Organic Plant Production & Agroecosystems Research in the Tropics and Subtropics, University of Kassel, Germany; Corresponding Author,

b. Formerly Architect, Urban planning Division, Kolkata Metropolitan Development Authority, Kolkata, India

c. Fellow, Institute of Town Planners and Associate, Indian Institute of Architects, India

Submitted: 8 January 2014; Revised 9 May 2014; Accepted for publication: 28 May 2014; Published: 10 June 2014

Abstract

The role urban and peri-urban agriculture (UPA) plays in reducing urban poverty and ensuring environmental sustainability was recognized by the Millennium Development Goals (MGDs). India is the world's largest democratic nation with a population of 1.2 billion. The rapid urbanization and high proportion of people below the poverty line along with higher migration to urban areas make India vulnerable to food crisis and urbanization of poverty. Ensuring jobs and food security among urban poor is a major challenge in India. The role of UPA can be well explained and understood in this context. This paper focuses on the current situation of UPA production in India with special attention to wastewater irrigation. This question is being posed about the various human health risks from wastewater irrigation which are faced by farmers and labourers, and, secondly by consumers. The possible health hazards involve microbial pathogens as well as helminth (intestinal parasites). Based on primary and secondary data, this paper attempts to confirm that UPA is one of the best options to address increasing urban food demand and can serve to complement rural supply chains and reduce ecological food prints in India. "Good practice urban and peri-urban agriculture" necessitates an integrated approach with suitable risk reduction mechanisms to improve the efficiency and safety of UPA production.

Keywords: *Health risks; Millennium Development Goals; Urban and peri-urban agriculture; Wastewater use*

Introduction

The significant role of urban and peri-urban agriculture (UPA) in the fulfilment of the Millennium Development Goals (MGDs), especially reducing urban poverty and hunger (MDG 1) and ensuring environmental sustainability (MGD 7), has been well recognized (Von Braun *et al.*, 2004; Mougeot, 2005). "Urban and peri-urban agriculture can

be broadly defined as the production, processing and distribution of foodstuff from crop and animal production, fish, and ornamental flowers within and around urban areas" (Mougot, 2000). UPA production systems were based on intensive and high input management practices on scarce lands (Smith *et al.*, 1996; Pearson *et al.*, 2010)




Figure 1 Social, ecological and economic dimensions and various types of urban and peri-urban agriculture (modified after; De Zeeuw et al., 2011).

depending on limited resources including water (Smit and Nasr, 1992). The achievement of food security can be asserted by increasing production, preventing post-harvest losses, improving the distribution network and enabling accessibility of food to poor people. The UPA can potentially fill the hunger gap by enhancing the access to and distribution of food in urban areas (Lee-Smith, 2010). Initially urban agriculture was started as a part of leisure time activity as well

as subsistence in world wars; a radical transformation of the objectives of UPA occurred during the early 1980s (Figure 1 and 2). In African countries especially Nigeria (Kano), Zimbabwe (Harare), and Tanzania (Dar-Es-Salaam) UPA became an integral part of the permanent landscape (Smith, 2001; Bryld, 2003). The driving force behind the transformation in UPA towards market oriented production is the increase in urban population




Figure 2 Benefits and services offered by various types of urban and peri-urban agriculture (modified after; Devenish, 2009).

and poverty (Cohen, 2006; Obuobie *et al.*, 2006; Hill *et al.*, 2007; Sinha, 2009; Ward, 2013).

UPA production systems may reduce the ecological foot print of cities and allow for synergies between urban domestic and industrial sectors (Jansen, 1992; Midmore and Jansen, 2003; De Zeeuw *et al.*, 2011). More than 800 million farmers are involved in UPA production, of which 200 million depend on wastewater for irrigation (UNDP, 1996; Bahri, 2009). The use of non-treated wastewater and industrial pollution make this system much more prone to high level heavy metals and microbial load thereby challenging the quality of urban produce (Diogo *et al.*, 2010; Abdu *et al.*, 2011a; Abdu *et al.*, 2011b; Kiba *et al.*, 2012; Safi and Buerkert, 2012).

Our objectives of the study were (1) to give an overview of the UPA production systems in India which are irrigated with wastewater (2) to compare the status quo of heavy metals present in different locations where UPA is practiced in India.

Methodology

The methodology used in this paper was based on primary and secondary data from various resources. In this paper, we review the current status of UPA production in India using wastewater as an irrigation source. We collected different research articles and books from multiple academic databases. Thus, this paper

built partially on our own research work as well as on a literature survey. The negative impacts on environment and health using wastewater were also mentioned. The results were analysed by

descriptive statistics using SPSS 12.0 (SPSS Inc., Chicago, IL, USA) and graphs were generated using Sigma plot 7 (Systat Software Inc., Chicago, IL, USA).




Figure 3 Population growth and Gross Domestic Product (GDP) growth in India from 1950 to the 2000s (after Panagariya, 2008).

Urbanization in India

India is the world's largest democratic nation with a population of 1.2 billion (Figure 3) where one third of the poor still lives below the poverty line of 1 US\$ per day (Datt and Ravallion, 2002; Deaton and Dreze, 2002; Census India, 2011). The current growth rate, genie coefficient and multidimensional poverty index of India is 4.0, 36.8 and 0.28 respectively (Krueger, 2008; Panagariya, 2008). In 1950, it was estimated that 82% of the total population lived in rural areas and agriculture contributed 56% of the Gross Domestic Product (GDP). While in 2010, due to rapid

urbanization the rural population declined to 70%, the urban population keeps increasing at a faster rate (Panagariya, 2008; Census India, 2011). The major populated metropolitan regions in India (Table 1) include the National Capital Region (NCR), Mumbai Metropolitan Region (MMR), Kolkata Metropolitan Region (KMR), Chennai Metropolitan Region (CMR), Bangalore Metropolitan Region (BMR) and Hyderabad Metropolitan Region (HMR). Based on the above facts, current population growth and relevance of UPA, our research question deals with the safety aspect

of wastewater irrigation in UPA production systems in India.

By 2025 India is expected to be the world's most populous country, thus bringing down the land to man ratio even further. The urban population will surpass the rural population in the course of the next decade. Land will become a more scarce resource for the farmers in the peri-urban areas and subsequently villages were transformed to urban areas. India's rate of urbanization is estimated to be about 3.5% per annum (Datt and Ravallion, 2002; Panagariya, 2008). The projection is that by 2020, about 50% of the total population of India will live in urban areas. Population explosion and the migration of people towards urban area create more pressure on food, shelter, water and basic necessities (Cohen, 2006). Migration from rural area to urban area is a common phenomenon in India, where people look for better employment, education, services and financial gain. Transformations in villages, alternative jobs in construction and various industries, poor productivity in agricultural labours seeking better job opportunities and climate change are some key factors triggering the decline of farming activities in rural and peri-urban areas (Sharma and Bhaduri, 2006; Martin, 2010).

It is estimated that Indian cities will generate 70% of the new jobs and 70% of Indian GDP in the year 2030 (McKinsey Global Institute, 2010). Employment and surging growth in Indian cities drove their population to 340 million in 2008 and could reach 590 million by 2030 (Panagariya, 2008). Poverty and a lack of gainful employment in rural areas drive people to the cities for

work and livelihood (Bhowmik, 2000). Five states in India will have more than 50% of urban population including Tamil Nadu, Gujarat, Maharashtra, Karnataka and Punjab by 2030 (McKinsey Global Institute, 2010).

UPA production in India

Urban agriculture in India is witnessing a beginning with piecemeal efforts in a few cities. Against the backdrop of tremendous population growth, haphazard and unplanned urbanization, growing food scarcity and increasing fruit and vegetable prices, there is the growing presence of urban agriculture in some form or other in every city. It is used as a resource conserving industry, where waste is converted to resource. It creates a diverse ecology where fruit trees, vegetable plants and fish production co-exist with the built environment of the urban poor, mostly migrants, making an ecologically sustainable scenario. The phenomena usually take place in the low-lying city fringes, which play important roles such as: a) controlling floods b) functioning as a workshop of 'resource conservation industry', where the open loop of thrown away garbage becomes a closed loop by converting it into resources c) supplying food and commodities to the city to keep the metabolism d) providing employment opportunities in the informal sector.

Urban agriculture can be considered one aspect for mitigating food insecurity and malnutrition among urban poor in India. In addition to livelihood opportunities, urban waste management also greatly improved (Gupta and Gangopadhyay, 2013). Usage of

wastewater in agricultural production systems can significantly provide an uninterrupted supply of resources, especially irrigation water and nutrients which can offer improved crop yields (Bahri, 1999; Kretschmer *et al.*, 2002; Bahri, 2009). The major UPA production centres in India were the six metropolitan cities (Figure 4 and 5).

Wastewater use

Due to the shortage and demand of fresh water, wastewater is often used as a valuable resource to meet the demand. It was estimated that on a global scale about 20 million ha (hectares) of land were irrigated with wastewater (Hamilton *et al.*, 2007). Wastewater reuse in agriculture a) provides an additional source of water, nutrient and organic matter b) reduces the discharge to the surface water c) improve the economic efficiency of investments in wastewater disposal and irrigation (Khouri *et al.*, 1994; Verma and Rakshit, 2010). Wastewater is comprised of domestic effluents, commercial establishments, industrial effluents and urban runoff. The constituents may vary spatially and temporally depending on the climatic conditions (Scott *et al.*, 2004). The most relevant wastewater usage includes a) direct use of untreated wastewater b) direct use of treated wastewater c) indirect use of wastewater (Figure 6 and 7). Use of wastewater for irrigation and aquaculture is considered a part of the informal sector in India and receives less attention from governments (Buechler *et al.*, 2002; Buechler and Devi, 2003; Buechler and Mekala, 2005; Amerasinghe *et al.*, 2013). The majority of

wastewater generated in India was taken from industrial effluents. According to Strauss and Blumenthal (1990), 73,000 ha were irrigated with wastewater in India. The majority of wastewater irrigation occurs along rivers adjacent to growing cities especially Delhi, Kolkata, Coimbatore, Hyderabad and Varanasi (Scott *et al.*, 2004).

Wastewater usage in Hyderabad

The seasonal flow pattern of the Musi River has been lost and most of the urban wastewaters drain into the river. Currently these wastewaters were used for growing para grass (*Urochloa mutica*) and paddy (*Oryza sativa*) where fodder grass targets the urban dairy market. Other wastewater irrigated crops include banana, coconut and vegetables with an estimated area of wastewater irrigation around 200 km² (Ensink *et al.*, 2005; Van Rooijen *et al.*, 2005). Along the Musi River it was estimated that 2,100 ha land were irrigated with wastewater to cultivate paddy (Mekala *et al.*, 2008; Srinivasan and Reddy, 2009). More than 13 vegetables were also grown under waste water irrigation which includes spinach (*Spinacia oleracea* L.), malabar spinach (*Basella Alba* L.), red amaranth (*Amaranthus cruentus* L.), mint (*Mentha spicata* L.), coriander (*Coriandrum sativum* L.), lady's finger/okra (*Abelmoschus esculentus* L.), taro (*Colocasia esculenta* L.) and common purslane (*Portulaca oleracea* L.). Flower production, especially jasmine (*Jasminum officinale* L.) using wastewater is also reported by Buechler *et al.* (2002).



Figure 4 Major UPA production systems from India: (top left), Railway Garden (RG) from the Mumbai Metropolitan Region; (top right), RG irrigated with wastewater; (middle left), farmers washing the white radish harvested from RG; (middle right), wastewater lagoon for fish production from Kolkata Metropolitan Area; (bottom left), mustard cultivation using wastewater in Kolkata Metropolitan Area; (bottom right), wastewater irrigation using motor pumps from lagoon in Kolkata Metropolitan Area.




Figure 5 Map of India showing the location of the UPA production areas (marked as star symbol). Source: own illustration.

Wastewater usage in Mumbai Metropolitan Region

In the Mumbai Metropolitan Region (MMR) the Indian Railways plays a major role in UPA production. Under the scheme "Grow more food", the Indian Railway companies has rented since 1975 unutilized land near railway tracks and stations to railway class IV employees and non-railway employees for promoting the cultivation of vegetables and as a part of corporate social responsibility (CSR). In the MMR, about 176 hectare of land were allotted among 282 railway employees who transformed this land to productive railway gardens by growing vegetables such as okra, spinach, red amaranth and taro which were predominantly irrigated with wastewater (Table 2; Vazhacharickal and Buerkert, 2012; Vazhacharickal et al., 2013).

Wastewater usage in Delhi Metropolitan Region

It was estimated that 1,700 ha of land irrigated with wastewater to grow cucurbits (*Cucurbita pepo*), eggplant (*Solanum melongena* L.), okra, coriander in summer and spinach, mustard (*Brassica juncea*), cauliflower (*Brassica oleracea* L.) and cabbage (*Brassica oleracea* var. *capitata*) in the winter across the Delhi metropolitan region (Mekala et al., 2008).

Wastewater usage in Kolkata Metropolitan Region

The east Kolkata sewage fisheries represent the largest single wastewater usage system in aquaculture in the world (Pescod, 1992). It was estimated that 12,800

tonnes of paddy, 6,900 tonnes of fish and 0.7 tonnes of vegetables were produced in this region (Chattopadhyay, 2002; Mukherjee et al., 2013).

Wastewater usage in Hubli-Dharwad

Wastewater irrigated agro-forestry systems were observed in villages near Hubli-Dharwad in Karnataka. The tree species include sapodilla (*Manilkara zapota*), guava (*Psidium guajava*), coconut (*Cocos nucifera* L.), mango (*Mangifera indica*), areca nut (*Areca catechu*) and teak (*Tectona grandis*). This complex agrosilivultural system also contains neem (*Azadirachta indica*), tamarind (*Tamarindus indica* L.), banana (*Musa acuminata*), pomegrate (*Punica granatum* L.) and mulberry (*Morus alba* L.; Bradford et al., 2003).

Wastewater usage in Kanpur

During 2004 it was estimated that 2,770 farmers were involved in wastewater agriculture stretching to 2,500 ha. About 400 million litres of wastewater were discharged in Kanpur per day (Gupta, 2013).

Wastewater usage in Coimbatore

In Coimbatore the major wastewater irrigation can be seen near Ukkadam with a sewage farm of size 35 ha cultivated with Guinea grass (*Megathyrsus maximus*), para grass, elephant grass (*Pennisetum purpureum*) and coconut (Jeyabaskaran and Sree Ramulu, 1996; Chitdeshwari et al., 2003; Somasundaram, 2003; Malarkodi et al., 2007).

Wastewater usage in Varanasi

In Varanasi it was estimated that 240 million litres of swages were generated. The major vegetables grown were radishes (*Raphanus sativus* L.), turnips (*Brassica rapa* var. *rapa* L.), carrots (*Daucus carota*), tomatoes (*Solanum lycopersicum* L.), cauliflower, eggplant, potatoes (*Solanum tuberosum* L.), cabbage, spinach and coriander which are targeted toward the urban market in Varanasi. The major villages using wastewater irrigation were Dinapur, Khalispur, Kotwa, Kamanli and Shiwar (Ghosh *et al.*, 2012).

Wastewater usage in Chennai

The major crops were millets (*Pennisetum glaucum* L.), sugarcane (*Saccharum* L.) and paddy, (Janakarajan *et al.*, 2010), but the usage of wastewater has not been officially reported by any authors, even though the usage with wastewater is common in peri-urban areas.

Health hazards associated with UPA production using wastewater

Human health risks from wastewater irrigation includes primarily farmers and labourers and secondly consumers. The possible health hazards involve microbial pathogens as well as helminth. In addition the soil and crop contamination of organic and inorganic trace elements were also reported in wastewater irrigation (Ganeshamurthy *et al.*, 2008). Farmers using wastewater irrigation are more prone to helminthes and skin problems than a famer using fresh water. (Trang *et al.*, 2007; Qadir *et al.*, 2010). Continuous

irrigation with wastewater may lead to heavy metal accumulation in the soil and their subsequent transfer to the vegetables (Table 3 and 4). Due to the persistent nature of these heavy metals, they may accumulate in vital organs and can cause numerous health disorders.

Concentrations of total Pb (lead) and Cd (cadmium) exceeded the safety thresholds (Table 6) in many vegetables, especially in spinach (3.8 and 1.8 mg kg⁻¹), green amaranth (3.3 and 0.2 mg kg⁻¹), white radish leaves (6.8 and 0.5 mg kg⁻¹), and white radish root (5.7 and 0.2 mg kg⁻¹). In all samples analysed Hg (mercury) and Ni (nickel) were below detection limit. The presence of Pb and Cd in plant material may in the long run create health hazards for the consumers. These metals are often accumulated in leafy vegetables and root crops when compared to fruits and seeds. The supply of nutrients in the irrigation water (Figure 8) may leads to ground water leaching which may affect the apparent nutrient usage efficiency of the system.

Management interventions for risk reduction

Human and environmental risk of wastewater usage can be reduced by a set of combined measures. These include a) water quality improvements b) human exposure control c) farm level wastewater and d) harvest and post-harvest interventions (WHO, 2006; Qadir *et al.*, 2010). Improvement in the quality of wastewater can be achieved by combination of primary and secondary treatment. Protective gadgets especially boots and gloves




Figure 6 Topology of wastewater usage (adapted from Keremane, 2007).

and changing irrigation methods can also lower the risk. Farm management practices including crop diversification irrigation and manure requirement also play a vital role. Finally the harvest and post-harvest interventions can also substantially reduce the health risk (Qadir *et al.*, 2010).

Government and public authorities lack the technical and management options to reduce the health risks associated with wastewater usage. Policies should be made to reduce the negative impacts of wastewater usage while supporting its benefits including the source as a valuable nutrient and organic matter. Management of wastewater and proper treatment will substantially reduce the possible

health risks. Conducting public awareness programmes will certainly make the farmers and consumers aware of the negative aspects of wastewater irrigation thereby encouraging them to adopt hygienic management practices.

Discussion

The contribution of UPA towards food security and employment opportunities was well appreciated in India. It is unquestionable that the UPA provides access of food to the urban poor at a cheap price. In addition to all these plus points, the usage of non-treated wastewater may impose long term health problem as well as the accumulation of heavy metals in the soils. Heavy metals were reported in various UPA production systems depending on




Figure 7 Dimensions of water usage (adapted from Scott *et al.*, 2004).

wastewater as irrigation sources. Heavy metal contaminations in vegetables grown using wastewater were reported in Hubli-Darwad (Hunshal *et al.*, 1997), Varanasi (Ghosh *et al.*, 2012), Hyderabad (Srikanth and Reddy, 1991) and Coimbatore (Malarkodi *et al.*, 2007). Consumption of vegetables with heavy metals may lead to accumulation and its long term use can pose serious health risks to the consumers. Attention is needed for the monitoring of industrial effluents, hygienic practices among farmers as well as the productive use of wastewater for irrigation. A permanent solution to prevent the heavy metals into the food chain seems to be less practicable in the

Indian scenario; however, the methods should be adopted to reduce the integration of these heavy metals, particularly through wastewater treatment as well as bioremediation methods.

Conclusions

Ensuring jobs and food security among the urban poor is a major challenge in underdeveloped and developing countries. Urban and peri-urban agriculture is one of the best options to address increasing urban food demand and can complement rural supply chains and reduce ecological foot prints.




Figure 8 Box plot showing the chemical characteristics of the irrigation water in ten UPA production systems across the Mumbai Metropolitan Region (India) and recommended threshold levels (FAO, 2010) displayed above the whisker.

The growing water demands and release of untreated wastewater pose a big challenge to environmental sustainability. Irrigation with wastewater is a reality and common practice in India. However, the possible health risks associated with them should receive attention from the policy makers and stakeholders. An integrated approach with suitable risk reduction mechanism would improve the efficiency and safety of these UPA production systems which can be called “Good practice urban and peri-urban agriculture”.

Acknowledgements

The first author would like to thank International Centre of Development and Decent Work (ICDD) at University of Kassel, Germany and Fiat Panis Foundation (Ulm, Germany) for providing a scholarship and necessary financial support. The authors also express their gratitude towards the journal editor, Linda Splinter and anonymous reviewers of “*Future of Food Journal*” for their support, critical comments and feedback.

References

- Abdu, N., Abdulkadir, A., Agbenin, J. O., & Buerkert, A. (2011a). Vertical distribution of heavy metals in wastewater-irrigated vegetable garden soils of three West African cities. *Nutrient Cycling in Agroecosystems*. 89(3): 387-397.
- Abdu, N., Agbenin, J. O., & Buerkert, A. (2011b). Phytoavailability, human risk assessment and transfer characteristics of cadmium and zinc contamination from urban gardens in Kano, Nigeria. *Journal of the Science of Food and Agriculture*. 91(15): 2722-2730.
- Amerasinghe, P., Bhardwaj, R. M., Scott, C., Jella, K., & Marshall, F. (2013). Urban wastewater and agricultural reuse challenges in India. Colombo, Sri Lanka: International Water Management Institute.
- Awasthi, S. K. (2000). Prevention of Food Adulteration Act No. 37 of 1954, Central and State rules as amended for 1999, 3rd Ed., New Delhi, India.
- Bahri, A. (1999). Agricultural reuse of wastewater and global water management. *Water Science and Technology*. 40(4): 339-346.
- Bahri, A. (2009). Managing the other side of the water cycle: Making wastewater an asset. Stockholm, Sweden: Global Water Partnership.
- Bhowmik, S. 2000. Hawkers and urban informal sector: A study of street vending in seven cities. National Alliance Street Vendors of India. Available from <http://www.nasvinet.org/userfiles/file/A%20study%20of%20street%20vending%20in%20seven%20cities.pdf> (Retrieved on August 02, 2013)
- Bradford, A., Brook, R., & Hunshal, C. S. (2003). Wastewater irrigation in Hubli-Dharwad, India: Implications for health and livelihoods. *Environment and Urbanization*. 15(2): 157-170.
- Bryld, E. (2003). Potentials, problems, and policy implications for urban agriculture in developing countries. *Agriculture and human values*. 20(1): 79-86.
- Buechler, S., & Mekala, G. D. (2005). Local responses to water resource degradation in India: Groundwater farmer innovations and the reversal of knowledge flows. *The Journal of Environment & Development*. 14(4): 410-438.
- Buechler, S., & Devi, G. (2003). Household food security and wastewater-dependent livelihood activities along the Musi River in Andhra Pradesh, India. Geneva, Switzerland: World Health Organization.

- Buechler, S., Devi, G., & Raschid, L. (2002, December). Livelihoods and wastewater irrigated agriculture: Musi River in Hyderabad city, Andhra Pradesh, India. *Urban Agriculture Magazine*, 14-17. Available from <http://www.ruaf.org/sites/default/files/Livelihoods%20and%20Wastewater%20Irrigated%20Agriculture.pdf> (Retrieved on May 18, 2013)
- CCME. (2001). Canadian water quality guidelines for the protection of aquatic life. Winnipeg, Canada: Canadian Council of Ministers of the Environment.
- Census India. (2011). Population of India. Retrieved from organizational web site: <http://censusindia.gov.in/>
- Chattopadhyay, K. (2002). Jalabhumir Kolkata. Kolkata, India: Indian Statistical Institute.
- Chitdeshwari, T., Duraisami, V. P., & Singh, M. V. (2003). Chemical composition of sewage effluents of major cities of Tamil Nadu. In: *Wastewater Treatment and Waste Management: Proceedings of the International Conference on Water and Environment (WE-2003)*, December 15-18, 2003, Bhopal, India (Vol. 2, p. 227).
- Cohen, B. (2006) Urbanization in developing countries: current trends, future projections, and key challenges for sustainability. *Technology in Society*. 28(1): 63-80.
- Datt, G., & Ravallion, M. (2002). *Is India's Economic Growth Leaving the Poor Behing?*. New York, NY: World Bank.
- Department of Water Affairs and Forestry. (1996). *South African Water Quality Guidelines (second edition). Volume 1: Domestic Use*. Pretoria, South Africa: Department of Water Affairs and Forestry.
- Devenish, C. (2009). *Urban Agriculture for Poverty Alleviation: A Case Study of Hyderabad, India*. Available from: <http://researcharchive.vuw.ac.nz/handle/10063/1113> (Retrieved on June 27, 2013)
- De Zeeuw, H., van Veenhuizen, R., & Dubbeling, M. (2011). The role of urban agriculture in building resilient cities in developing countries. *The Journal of Agricultural Science*. 149(S1): 153-163.
- Deaton, A., & Dreze, J. (2002). Poverty and inequality in India: a re-examination. *Economic and Political Weekly*. 37(36): 29-3748.
- Diogo, R. V., Buerkert, A., & Schlecht, E. (2010). Horizontal nutrient fluxes and food safety in urban and peri-urban vegetable and millet cultivation of Niamey, Niger. *Nutrient cycling in agroecosystems*. 87(1): 81-102.

- Ensink, J. H., van der Hoek, W., Mukhtar, M., Tahir, Z., & Amerasinghe, F. P. (2005). High risk of hookworm infection among wastewater farmers in Pakistan. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 99(11), 809-818.
- FAO/WHO. (2001). Food additives and contaminants. Joint Codex Alimentarius Commission. New York, USA: Food and Agriculture Organization of the United Nations/ World Health Organization.
- FAO. (2010). Irrigation water quality guidelines. New York, USA: Food and Agriculture Organization.
- Ganeshamurthy, A. N., Varalakshmi, L. R., & Sumangala, H. P. (2008). Environmental risks associated with heavy metal contamination in soil, water and plants in urban and periurban agriculture. *Journal of Horticultural Sciences*. 3(1): 1-29.
- Ghosh, A. K., Bhatt, M. A., & Agrawal, H. P. (2012). Effect of long-term application of treated sewage water on heavy metal accumulation in vegetables grown in Northern India. *Environmental monitoring and assessment*. 184(2): 1025-1036.
- Gupta, K. (2013). Water quality, wastewater generation and sewerage system in urban areas: a case study of Kanpur city. *Indian Streams Research Journal*. 3(9): 1-6.
- Gupta, R., & Gangopadhyay, S. G. (2013). Urban Food Security through Urban Agriculture and Waste Recycling: Some Lessons for India. *VIKALPA*. 38(3): 13.
- Hamilton, A. J., Stagnitti, F., Xiong, X., Kreidl, S. L., Benke, K. K., & Maher, P. (2007). Wastewater irrigation: The state of play. *Vadose Zone Journal*. 6(4): 823-840.
- Hill, K., Quinnelly, D.D., & Kazmierowski, K. (2007). Urban agriculture in Naga city. Cultivating sustainable livelihoods. Available from organizational web site:
<http://www.cityfarmer.org/Urban%20Agriculture%20Group%20Final%20Report.pdf> (Retrieved on July 03, 2013)
- Hunshal, C. S., Salakinkop, S. R., & Brook, R. M. (1997). Sewage irrigated vegetable production systems around Hubli-Dharwad, Karnataka, India'. *The Kasetart Journal (Natural Sciences)*. 32(32): 1-8.
- Janakarajan, S., Butterworth, J., Moriarty, P., & Batchelor, C. (2010). Strengthened city, marginalised peri-urban villages: stakeholder dialogues for inclusive urbanisation in Chennai, India. In: *Peri-Urban Water Conflicts Supporting dialogue and negotiation*. Delft: International Water and Sanitation Centre.

- Jansen, H.G.P. (1992). Supply and demand of AVRDC mandate crops in Asia: implications of past trends for future developments. Tainan, Taiwan: Asian Vegetable Research and Development Centre.
- Jeyabaskaran, K. J., & Sree Ramulu, U. S. (1996). Distribution of heavy metals in soils of various sewage farms in Tamil Nadu. *Journal of the Indian Society of Soil Science*. 44(3): 401-404.
- Keremane, G. (2007). Urban wastewater reuse, an alternative source for agricultural irrigation-A review. Available from organizational web site: http://www.soil.tubs.de/lehre/Master.Irrigation/2011/Lit/1_UrbanWastewaterReuse.8189KERRev.pdf (Retrieved on May 24, 2013)
- Khoury, N., Kalbermatten, J. M., & Bartone, C. (1994). The reuse of wastewater in agriculture: A guide for planners. UNDP-World Bank Water and Sanitation Program, Washington, DC: World Bank.
- Kiba, D. I., Zongo, N. A., Lompo, F., Jansa, J., Compaore, E., Sedogo, P. M., & Frossard, E. (2012). The diversity of fertilization practices affects soil and crop quality in urban vegetable sites of Burkina Faso. *European Journal of Agronomy*. 38: 12-21.
- KMDA. 2012. Functioning of six metropolitan development authorities in India: a comparative analysis. Kolkata Metropolitan Development Authority, Kolkata, India.
- Kretschmer, N., Ribbe, L., & Gaese, H. (2002). Wastewater reuse for agriculture. *Technology Resource Management & Development-Scientific Contributions for Sustainable Development*. 2: 37-64.
- Krueger, A. O. (2008). The Role of Trade and International Economic Policy in Indian Economic Performance. *Asian Economic Policy Review*. 3(2): 266-285.
- Lee-Smith, D. (2010). Cities feeding people: an update on urban agriculture in equatorial Africa. *Environment and Urbanization*. 22(2): 483-499.
- Malarkodi, M., Krishnasamy, R., Kumaraperumal, R., & Chitdeshwari, T. (2007). Characterization of heavy metal contaminated soils of Coimbatore district in Tamil Nadu. *Journal of Agronomy*. 6(1): 147.
- Martin, P. 2010. Climate change, agricultural development, and migration. The German Marshall Fund of the United States. Available from http://www.gmfus.org/galleries/default-file/PMartin_V2.pdf (Retrieved on August 12, 2013)
- McKinsey Global Institute. 2010. India's urban awakening: Building inclusive cities, sustaining economic growth. McKinsey & Company, Mumbai, India.

- Mekala, G. D., Samad, B., & MadarBoland, A. M. (2008). Wastewater reuse and recycling systems: a perspective into India and Australia. Colombo, Sri Lanka: International Water Management Institute.
- Midmore, D.J., & Jansen, H.G.P. (2003). Supplying vegetables to Asian cities: is there a case of peri-urban production? *Food Policy*. 28: 13-27.
- Mougeot, L.J. (2005). *Agropolis: The Social, Political and Environmental Dimensions of Urban Agriculture*. London, UK: Earthscan.
- Mougeot, L. J. (2000). Urban agriculture: definition, presence, potentials and risks. *Growing cities, growing food: Urban agriculture on the policy agenda*, DSE, Germany. 1-42.
- Mukherjee, V., Das, A., Akhaand, A., & Gupta, G. (2013). Toxicity and Profitability of Rice Cultivation under Waste-Water Irrigation: The Case of the East Calcutta Wetlands. *Ecological Economic*. 93: 292-300.
- Obuobie, E., Keraita, B., Danso, G., Amoah, P., Cofie, O.O., Raschid-Sally, L., & Drechsel, P. (2006). *Irrigated urban vegetable production in Ghana: Characteristics, benefits and risks*. Accra, Ghana: International Water management Institute.
- Panagariya, A. (2008). *India: The emerging giant*. New York, USA: Oxford University Press.
- Pearson, L. J., Pearson, L. & Pearson, C. J. (2010) Sustainable urban agriculture: stocktake and opportunities. *International Journal of Agricultural Sustainability*. 8(1-2): 7-19.
- Pescod, M. B. (1992). *Wastewater treatment and use in agriculture*. FAO Irrigation and Drainage Paper 47, Rome: Food and Agriculture Organization.
- Qadir, M., Wichelns, D., Raschid-Sally, L., McCornick, P. G., Drechsel, P., Bahri, A., & Minhas, P. S. (2010). The challenges of wastewater irrigation in developing countries. *Agricultural Water Management*. 97(4): 561-568.
- Safi, Z., & Buerkert, A. (2012). Heavy metal and microbial loads in sewage irrigated vegetables of Kabul, Afghanistan. *Journal of Agriculture and Rural Development in the Tropics and Subtropics*. 112(1): 29-36.
- Scott, C. A., Faruqui, N. I., & Raschid-Sally, L. (2004). 1. Wastewater Use in Irrigated Agriculture: Management Challenges in Developing Countries. In: *Wastewater use in irrigated agriculture: confronting the livelihood and environmental realities*. Oxfordshire: CABI Publishing.

- Sharma, A., Bhaduri, A. 2006. The "Tipping Point" in Indian agriculture: Understanding the withdrawal of the Indian rural youth. Draft prepared for the IWMI-CPWF project on "Strategic Analysis of National River Linking Project of India". Available from <http://www.iwmi.cgiar.org/Publications/Other/PDF/Paper%207%20of%20NRLP%20series%201.pdf> (Retrieved on April 21, 2013)
- Sinha, A. (2009). Agriculture and Food Security: Crises and Challenges Today. *Social Action*. 59(2): 1-16.
- Smit, J., Nasr, J., & Ratta, A. (1996) *Urban Agriculture: Food, Jobs and Sustainable Cities*. New York, NY: The Urban Agriculture Network Inc.
- Smit, J., & Nasr, J. (1992). Urban agriculture for sustainable cities: using wastes and idle land and water bodies as resources. *Environment and Urbanization*. 4(2): 141-152.
- Smith, O. B. (2001). *Overview of urban agriculture in Western African cities*. Ottawa, Canada: International Development Research Centre.
- Somasundaram, J. (2003, November 6). Imbibing toxic heavy metals through leafy vegetables. *The Hindu*. Available from <http://www.hindu.com/seta/2003/11/06/stories/2003110600110300.htm> (Retrieved on July 15, 2013)
- Srikanth, R., & Reddy, S. (1991). Lead, cadmium and chromium levels in vegetables grown in urban sewage sludge-Hyderabad, India. *Food chemistry*. 40(2): 229-234.
- Srinivasan, J. T., & Reddy, V. R. (2009). Impact of irrigation water quality on human health: A case study in India. *Ecological Economics*. 68(11): 2800-2807.
- Strauss, M., & Blumenthal, U. J. (1990). Use of human wastes in agriculture and aquaculture; utilization practices and health perspectives. IRCWD report, 8.
- Trang, D. T., Van Der Hoek, W., Tuan, N. D., Cam, P. D., Viet, V. H., Luu, D. D., Konradsen, F., & Dalsgaard, A. (2007). Skin disease among farmers using wastewater in rice cultivation in Nam Dinh, Vietnam. *Tropical Medicine & International Health*. 12(s2): 51-58.
- UNDP (United Nations Development Program). 1996. *Urban Agriculture: Food, Jobs and Sustainable Cities*. Publication Series for Habitat II, Volume One. New York.
- Van Rooijen, D. J., Turrall, H., & Wade Biggs, T. (2005). Sponge city: water balance of mega-city water use and wastewater use in Hyderabad, India. *Irrigation and drainage*. 54(S1): S81-S91.

- Vazhacharickal, P. J., & Buerkert, A. (2012). Sustainable cities: an overview of the urban and peri-urban agricultural production in Mumbai Metropolitan Region (MMR). *Leituras de Economia Política*. 19: 69-87.
- Vazhacharickal, P. J., Predotova, M., Chandrasekharam, D., Bhowmik, S., & Buerkert, A. (2013). Urban and peri-urban agricultural production along railway tracks: a case study from the Mumbai Metropolitan Region. *Journal of Agriculture and Rural Development in the Tropics and Subtropics*. 114(2): 145-157.
- Verma, D. K., & Rakshit, A. (2010). Wastewater: a resource for agriculture. *International Journal of Agriculture, Environment and Biotechnology*. 3(3): 339-341.
- Von Braun, J., Swaminathan, M.S., & Rosegrant, M. W. (2004). Agriculture, food security, nutrition and the Millennium Development Goals., Essay in 2003-2004 IFPRI annual report, International Food Policy Research Institute. Available from <http://www.ifpri.org/sites/default/files/publications/ar03e.pdf>. (Retrieved on May 05, 2013)
- Ward, C. (2013). Urban agriculture helps to combat hunger in Indian's slums. World Watch Institute. Available from <http://www.worldwatch.org/urban-agriculture-helps-combat-hunger-india%E2%80%99s-slums-1> (Retrieved on June 10, 2013)
- WHO. (2006). WHO Guidelines for the Safe Use of Wasterwater Excreta and Greywater. New York, USA: World Health Organization.

Appendix

Rank	Metropolitan area	State/Territory	Governing authority	Area (km ²)	Population	Population density (km ²)
1	National Capital Region (NCR)	Delhi, Uttar Pradesh, Haryana	Delhi Development Authority (DDA)	1,483	21,753,000	9,340
2	Mumbai Metropolitan Region (MMR)	Maharashtra	Mumbai Metropolitan Region Development Authority (MMRDA)	4,355	20,748,000	4,764
3	Kolkata Metropolitan Area (KMA)	West Bengal	Kolkata Metropolitan Development Authority (KMDA)	1,886	14,617,000	12,883
4	Chennai Metropolitan Area (CMA)	Tamil Nadu	Chennai Metropolitan Development Authority (CMDA)	1,189	8,728,000	5,921
5	Bangalore Metropolitan Area (BMA)	Karnataka	Bangalore Development Authority (BDA)	1,276	9,645,000	7,600
6	Hyderabad Metropolitan Area (HMA)	Andhra Pradesh	Hyderabad Metropolitan Development Authority (HMDA)	7,100	7,749,000	7,826

Table 1 Major metropolitan areas in India and their characteristics during the year 2011 (KMDA, 2012).

Serial No	Common name (English)	Local name (Hindi)	Botanical name
1	Lady's finger/Okra	Bhindi	<i>Abelmoschus esculentus</i> L
2	Spinach	Palak	<i>Spinacia oleracea</i> L
3	Red amaranth	Lal Maat	<i>Amaranthus cruentus</i> L
4	Fenugreek	Methi	<i>Trigonella foenum-graecum</i> L
5	White radish	Mula	<i>Rhaphanus sativus</i> var. <i>longipinnatus</i>
6	Malabar spinach	Mayalu	<i>Basella alba</i> L
7	Green amaranth	Chawli	<i>Amaranthus tritris</i>
8	Sorrel leaves	Ambaadi	<i>Hibiscus sabdariffa</i> L
9	Taro	Alu	<i>Colocasia esculenta</i> L
10	Dill	Shepu	<i>Anethum graveolens</i> L

Table 2 Vegetables cultivated in UPA railway gardens of the Mumbai Metropolitan Region, India (Vazhacharickal et al., 2013).

Heavy metal *	Mean	SD	Min	Max	Thresholds	
Lead (Pb)	0.03	0.02	0.00	0.06	0.2 †a	0.05 ‡b
Cadmium (Cd)	0.38	1.08	0.00	3.47	0.05 †a	0.003 ‡b
Mercury (Hg)	0.003	0.001	0.001	0.005	0.002 †b	0.006 ‡b

* All values in mg l⁻¹

a Agricultural Standards

b Domestic Standards

† Department of Water Affairs and Forestry (1996), South African Water Quality Guidelines

‡ FAO (2010)

Table 3 Heavy metal concentrations in irrigation water collected from 10 UPA production systems in the Mumbai Metropolitan Region and recommended threshold levels.

Data	Cd	Pb	Cu	Zn	Cr	Ni
	(mg kg ⁻¹)					
Mean	0.4	3.8	146	133	429	135
SD	0.4	1.4	54	21	209	72
Max	1.1	6.9	231	161	791	247
Min	0.1	3.3	60	103	114	1.1
Thresholds						
India†	3-6	250-500	135-270	300	na	75-150
EU‡	3	300	140	600	150	20-100
UK‡	3	300	80-200	300	400	50-110
USA‡	20	150	170	1400	na	210

Table 4 Total cadmium (Cd), lead (Pb), copper (Cu), zinc (Zn), chromium (Cr) and nickel (Ni) concentrations in the surface soil (0-20 cm) of seven UPA production systems across the Mumbai Metropolitan Region, India.

Locations	Zn	Cu	Cd	Pb
	(mg kg ⁻¹)			
Bangalore	71.8	3.52	0.35	35.2
Kolkata	1300	160	4.0	170
Varanasi	87.9	33.5	2.7	18.3
Coimbatore	397.4	157.1	8.1	175.5
Hyderabad	2.9	4.3	0.4	8.1
PFA standard	300-600	135-270	3-6	25-50

Table 5 Total heavy metal content in soils of various UPA production system across India (modified after; Ganeshamurthy *et al.*, 2008).

Data	Cu	Zn	Cr	Pb	Cd
	(mg kg ⁻¹)				
Mean	7	40	4	5	0.6
SD	4	34	4	2	0.7
Max	13	96	11	7	1.8
Min	3	9	1	3	0.1
Safety thresholds					
WHO†	40	50	na	0.3	0.2
India‡	30	50	20	2.5	1.5

Table 6 Concentrations of total copper (Cu), zinc (Zn), chromium (Cr), lead (Pb) and cadmium (Cd) in green amaranth, spinach, white radish and paddy (mg kg⁻¹ dry weight) from UPA production systems (n=4) across the Mumbai Metropolitan Region, India.